

Journal Pre-proof

Clinical manifestations and outcome of SARS-CoV-2 infection during pregnancy

Yangli Liu , Haihong Chen , Kejing Tang , Yubiao Guo

PII: S0163-4453(20)30109-2
DOI: <https://doi.org/10.1016/j.jinf.2020.02.028>
Reference: YJINF 4471



To appear in: *Journal of Infection*

Accepted date: 27 February 2020

Please cite this article as: Yangli Liu , Haihong Chen , Kejing Tang , Yubiao Guo , Clinical manifestations and outcome of SARS-CoV-2 infection during pregnancy, *Journal of Infection* (2020), doi: <https://doi.org/10.1016/j.jinf.2020.02.028>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd on behalf of The British Infection Association.

Clinical manifestations and outcome of SARS-CoV-2 infection during pregnancy

Yangli Liu[#], Haihong Chen[#], Kejing Tang, Yubiao Guo*

The First Affiliated Hospital of Sun Yat-sen University, Guangzhou 510080, Province
Guangdong, P.R. China

#: These authors contributed equally to this work.

* This is corresponding author:

Yubiao Guo: Division of Pulmonary and Critical Care Medicine ,the First Affiliated
Hospital of Sun Yat-sen University, Guangzhou, Guangdong 510080, China.
yubiao_guo@163.com

Descargado para Anonymous User (n/a) en Cons
Para uso personal exclusivamente. No se permiten otros usos sin autorizac

**Keywords: Severe acute respiratory syndrome corona virus 2; 2019 novel
coronavirus disease; Clinical manifestations ;Pregnancy outcome**

Dear editors,

Tang and colleagues, in this Journal, drew readers attention to emerging COVID19[1].

We focused on the pregnant COVID19 patients. Given the maternal physiologic and immune function changes in pregnancy[2], pregnant individuals might face greater risk of getting infected by SARS-CoV-2 and might have more complicated clinical events. We described epidemiological, clinical characteristics, pregnancy and perinatal outcomes of all hospitalized pregnant patients diagnosed with COVID-19 in China.

We identified all hospitalized pregnant patients with laboratory-confirmed SARS-CoV-2 infection between December 8, 2019, and February 25, 2020 officially reported by the central government, in areas outside Wuhan, China. Information including age, geographic location, epidemiological history, prenatal course, maternal and newborn hospital course, discharge data and outcome were obtained by Centers for Disease Control and Prevention and Local Health Commission. When necessary, we attempted to contact local hospital or patients by telephone to supply missing information. This investigation was part of an emergency public health outbreak investigation and therefore not subject to institutional review board.

There were a total of 13 Chinese patients with SARS-CoV-2 admitted to hospitals outside of Wuhan (Table). There were 3 patients from Zhejiang, 3 from other cities of Hubei and 1 each from Fujian, Shanxi, Beijing, Guangdong, Jiangxi, Heilongjiang and Anhui. The maternal age ranged between 22 to 36 years. Two women were less than 28 weeks of gestation and the other 11 patients were in their

third trimesters at presentation. None of the patients had underlying medical disease.

Ten patients (77%) presented with fever (range 37.3–39.0°C), mostly accompanied with fatigue. Only 3 (23%) pregnant patients complained with dyspnea. 1 had no symptoms but got a positive RNA test result of oropharyngeal swabs after close contact to a diagnosed family member. 12 patients (92%) had a clear epidemiologic history, either with other family members affected or with linkage to Wuhan (residing in or visiting Wuhan or contact with visitors from Wuhan ≤ 2 weeks before the onset of infection).

Three of the patients (23%) improved after hospitalization and got discharged with an uncomplicated ongoing pregnancy. The other 10 patients (77%) all underwent caesarean section. Five of the 10 patients were delivered by emergency cesarean section because of pregnancy complications including fetal distress (in three of ten patients), premature rupture of the membrane (in one of ten) and stillbirth (in one of ten). Six patients (46%) had preterm labour between 32- 36 weeks of gestation.

Patient 6's condition deteriorated during hospitalization, prompting intensive care unit (ICU) admission with multiple organ dysfunction syndrome (MODS) including acute respiratory distress syndrome (ARDS) requiring intubation and mechanical ventilation, acute hepatic failure, acute renal failure and septic shock. As of February 25th, Patient 6 was still in the support of Extracorporeal Membrane Oxygenation (ECMO). The other 12 pregnant patients were all discharged with no obvious complication. Except for 1 stillbirth, nine newborn infants got a 1-min Apgar score of 10. There was no clinical or serologic evidence suggestive of vertical transmission of

SARS-CoV-2.

Previous studies suggested that COVID-19 is more likely to affect older males with comorbidities[3]. We reported 13 pregnant COVID-19 patients in China, indicating pregnant women also susceptible to SARS-CoV-2. Clinical manifestations of the pregnant COVID-19 patients in this study varied widely from asymptomatic to very severe, similar to previous report in non-pregnant patients [4]. Most of the pregnant patients had mild to moderate symptoms. Fever and fatigue were the principal symptoms, and less common symptoms were sore throat and shortness of breath. Almost all the patients had a clear epidemiologic history.

One of the 13 patients (7.6%) developed severe pneumonia requiring ICU care with multiple organ dysfunction syndrome in the third trimester in our study, similarity with the general population reported to be with critical rate of 5% [5]. Cytokine storm might be the reason for very severe cases since Chaolin Huang et al[6] found that compared with non-ICU patients, ICU patients had higher plasma levels of various cytokines.

Five patients of thirteen (38%) were delivered by emergency cesarean section because of pregnancy complications including fetal distress, premature rupture of the membrane and stillbirth. Six patients (46%) had preterm labour. These perinatal complications could be ascribed to the virus infection as well as the physiologic changes that reducing the woman intolerant to hypoxia during late pregnancy[7]. Fortunately, no severe neonatal asphyxia was observed in the nine livebirths and no vertical transmission was found.

In conclusion, our report showed pregnant women are also susceptible to SARS-CoV-2 infection. SARS-CoV-2 may increase health risks to both mothers and infants during pregnancy. Efforts should be taken to reduce the infection rate of SARS-CoV-2 both in pregnant and perinatal period, and more intensive attention should be paid to pregnant patients.

Conflict of interest

All authors declare that they have no competing interests.

References:

1. Tang JW, Tambyah PA, DSC H. Emergence of a novel coronavirus causing respiratory illness from Wuhan, China. *J Infect* **2020**;80:350-71.
2. Jamieson DJ, Theiler RN, Rasmussen SA. Emerging infections and pregnancy. *Emerg Infect Dis* **2006**;12:1638-43.
3. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* **2020**;395:507-13.
4. Xu XW, Wu XX, Jiang XG, et al. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series. *BMJ* **2020**;368:m606.
by Descargado para Anonymous User (n/a) en Cons
Para uso personal exclusivamente. No se permiten otros usos sin autorizac
5. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA* **2020**.
6. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **2020**;395:497-506.
7. Assiri A, Abedi GR, Al MM, Bin SA, Gerber SI, Watson JT. Middle East Respiratory Syndrome Coronavirus Infection During Pregnancy: A Report of 5 Cases From Saudi Arabia. *Clin Infect Dis* **2016**;63:951-3.

Table. Characteristics of 13 Hospitalized pregnant patients Infected With SARS-CoV-2

charac	Pati	Pati	Pati	Pati	Pati	Pati	Pati	Pati	Pati	Pati	Pati	Pati	Pati
teristic	ent	ent	ent	ent	ent	ent	ent	ent	ent	ent	ent	ent	ent
s	1	2	3	4	5	6	7	8	9	10	11	12	13
	Second trimester			> 28 wks of gestation									
Patient	28	24	33	29	35	31	30	36	26	32	30	22	30
age,y													
Gestati	25w	27w	32w	33w	34w	34w	35w	35+	36w	36+	37	38w	38+
onal						+		5		4			3
age at													
illness													
onset,													
wk													
Sympt	Fev	Fev	Pers	Fev	Fev	Fev	Dys	Fev	Fev	Fev	Fev	No	Fev
oms at	er,	er,	isten	er,	er	er,	pne	er,	er,	er,	er,		er,
onset	fatig	pea	t	pea		sore	a	cou	fatig	pea	fatig		pea
	ue	king	coug	king		thor		gh,	ue	king	ue		king
		at	h	at		at		dysp		at			at
		38.5		38				nea		39°C			37.3
		°C		°C									°C
		dys											
		pne								fatig			

Descargado para Anonymous User (n/a) en Cons
Para uso personal exclusivamente. No se permiten otros usos sin autorizac

	a				ue															
Epidemiology																				
history	Yes	No	Yes	No	No	No	No	No	No	No	Yes	Yes	No	Yes	Yes					
Other family members affected																				
Linkage to Wuhan																				
Compliance	No	No	No	No	PR	MO	Feta	No	No	Feta	Feta	No	No							
Locations					OM	DS	I			I	I									
Methylation	NA	NA	C-se	NA	C-se	C-se	C-se	C-se	C-se	C-se	C-se	C-se	C-se	C-se						
Method of			ction		ction	ction	ction	ction	ction	ction	ction	ction	ction	ction						

Descargado para Anonymous User (n/a) en Cons
Para uso personal exclusivamente. No se permiten otros usos sin autorizac

deliver

y

Maternal	Surv	Surv	Surv	Surv	Surv	Surv	Surv	Surv	Surv	Surv	Surv	Surv	Surv
Outcome	Survived	Survived	Survived	Survived	Survived	Survived	Survived	Survived	Survived	Survived	Survived	Survived	Survived

me

Premature	NA	NA	Yes	NA	Yes	Yes	Yes	Yes	Yes	No	No	No	No
------------------	----	----	-----	----	-----	-----	-----	-----	-----	----	----	----	----

ture

deliver

y

Descargado para Anonymous User (n/a) en Cons
Para uso personal exclusivamente. No se permiten otros usos sin autorizac

Fetal	NA	NA	10	NA	10	0	10	10	10	10	10	10	10
--------------	----	----	----	----	----	---	----	----	----	----	----	----	----

Apgar

score

Fetal	Surv	Surv	Surv	Surv	Surv	Died	Surv	Surv	Surv	Surv	Surv	Surv	Surv
Outcome	Survived	Survived	Survived	Survived	Survived		Survived	Survived	Survived	Survived	Survived	Survived	Survived

me

Vertical	NA	NA	No	NA	No	No	No	No	No	No	No	No	No
-----------------	----	----	----	----	----	----	----	----	----	----	----	----	----

I

transm

ission

Abbreviation: SARS-CoV-2=severe acute respiratory syndrome corona virus 2; PROM=premature rupture of membrane; MODS=multiple organ dysfunction syndrome; C-section=caesarean section